1467 (NH); St. Lucia, outside crocodile farm, 21 Aug. 1991, Hutchings & Hutchings 3017 (MO); grassland around Mt. Tabor, St. Lucia, Eastern Shores State Forest, 25 June 1989, Phillipson 3196 (MO, RUH); St. Lucia Estuary, dune forest, Aug. 1984, Balkwill & Cadman 1728 (NU); Mapalane, 16 Aug. 1991, Hutchings & Hutchings 3016 (MO); Kwambonambi, Amangwe Forest, 25 Aug. 1991, Hutchings & Hutchings 3019 (MO); Monzi turnoff on road to Mtubatuba, forest track, 9 Aug. 1977, Pooley 1870 (NU); Richards Bay, 26 July 1974, Ward 5 (NU), 4 Sep. 1970, Strey 9906 (NH).

KEY TO THE SUBSPECIES

1a. Flowers pink to red usually with dark red markings on the lower tepals; perianth tube 19-28

- (-32) mm long; dorsal tepal $11-15 \times 5$ mm subsp. laxa
- 1b. Flowers pale blue to whitish with dark blue to violet markings on the lower tepals; perianth tube (25-)27-32 mm long; dorsal tepal 9-11 × 4 mm subsp. azurea

Literature Cited

- Goldblatt, P. 1972. A revision of the genera *Lapeirousia* Pourret and *Anomatheca* Ker in the winter rainfall region of South Africa. Contr. Bolus Herb. 4: 1-111.
- logeny and biology of the African Iridaceae. Contr. Bolus Herb. 13: 1-74.